

# UCLA

## UCLA Previously Published Works

### Title

Impact of oral hygiene on head and neck cancer risk in a Chinese population

### Permalink

<https://escholarship.org/uc/item/9h94j423>

### Journal

Head & Neck, 39(12)

### ISSN

1043-3074

### Authors

Kawakita, Daisuke  
Lee, Yuan-Chin Amy  
Li, Qian  
et al.

### Publication Date

2017-12-01

### DOI

10.1002/hed.24929

Peer reviewed



Published in final edited form as:

Head Neck. 2017 December ; 39(12): 2549–2557. doi:10.1002/hed.24929.

## The impact of oral hygiene on head and neck cancer risk in a Chinese Population

Daisuke Kawakita, MD<sup>1,2,3</sup>, Yuan-Chin Amy Lee, PhD<sup>1</sup>, Qian Li, PhD<sup>4</sup>, Yuji Chen, BS<sup>1</sup>, Chien-Jen Chen, ScD<sup>5,6</sup>, Wan-Lun Hsu, PhD<sup>5</sup>, Pei-Jen Lou, MD, PhD<sup>7</sup>, Cairong Zhu, PhD<sup>8</sup>, Jian Pan, MD<sup>9</sup>, Hongbing Shen, MD, PhD<sup>10</sup>, Hongxia Ma, MD, PhD<sup>10</sup>, Lin Cai, MD, PhD<sup>11</sup>, Baochang He, PhD<sup>11</sup>, Yu Wang, MD<sup>12</sup>, Xiaoyan Zhou, MD<sup>13</sup>, Qinghai Ji, MD<sup>12</sup>, Baosen Zhou, MD, PhD<sup>14</sup>, Wei Wu, PhD<sup>14</sup>, Jie Ma, MD<sup>15</sup>, Paolo Boffetta, MD, MPH<sup>16</sup>, Zuo-Feng Zhang, MD, PhD<sup>17</sup>, Min Dai, PhD<sup>18</sup>, and Mia Hashibe, PhD<sup>1</sup>

<sup>1</sup>Division of Public Health, Department of Family and Preventive Medicine, and Huntsman Cancer Institute, University of Utah School of Medicine, Salt Lake City, UT, USA

<sup>2</sup>Department of Otorhinolaryngology, Head and Neck Surgery, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan

<sup>3</sup>Division of Molecular Medicine, Aichi Cancer Center Research Institute, Nagoya, Japan

<sup>4</sup>Departments of Preventive Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA

<sup>5</sup>Genomics Research Center, Academia Sinica, Taipei, Taiwan

<sup>6</sup>Graduate Institute of Epidemiology and Preventive Medicine, National Taiwan University, Taipei, Taiwan

<sup>7</sup>Department of Otolaryngology, National Taiwan University Hospital, Taipei, Taiwan

<sup>8</sup>Department of Epidemiology and Biostatistics, West China Center of Medical Sciences, Sichuan University, Sichuan, China

<sup>9</sup>Department of Oral Surgery, West China Hospital of Stomatology, Sichuan University, Sichuan, China

---

**Corresponding author:** Mia Hashibe, PhD, Division of Public Health, Department of Family and Preventive Medicine, Huntsman Cancer Institute, University of Utah School of Medicine, 2000 Circle of Hope, Salt Lake City, UT 84108, USA. mia.hashibe@utah.edu.

### Authorship

#### *Conception and design or analysis and interpretation of data*

Daisuke Kawakita, Yuan-Chin Amy Lee, Mia Hashibe

#### *Drafting of the manuscript or revising it for important intellectual content*

Daisuke Kawakita, Yuan-Chin Amy Lee, Mia Hashibe

#### *Final approval of the version to be published*

Qian Li, Yuji Chen, Chien-Jen Chen, Wan-Lun Hsu, Pei-Jen Lou, Cairong Zhu, Jian Pan, Hongbing Shen, Hongxia Ma, Lin Cai, Baochang He, Yu Wang, Xiaoyan Zhou, Qinghai Ji, Baosen Zhou, Wei Wu, Jie Ma, Paolo Boffetta, Zuo-Feng Zhang, Min Dai

### Conflict of interest statement

The authors declare no potential conflicts of interest.

<sup>10</sup>Department of Epidemiology and Biostatistics, Jiangsu key lab of Cancer Biomarkers, Prevention and Treatment, Collaborative Innovation Center for Cancer Personalized Medicine, School of Public Health, Nanjing Medical University, Nanjing, China

<sup>11</sup>Department of Epidemiology and Biostatistics, School of Public Health, Fujian Medical University, Fuzhou, China

<sup>12</sup>Department of Head and Neck Surgery, Fudan University Shanghai Cancer Center and Department of Oncology, Shanghai Medical College, Fudan University, Shanghai, China

<sup>13</sup>Department of Pathology, Fudan University Shanghai Cancer Center, Shanghai, China; Department of Oncology, Shanghai Medical College, Fudan University, Shanghai, China

<sup>14</sup>Department of Epidemiology, School of Public Health, China Medical University, Liaoning, China

<sup>15</sup>Department of Head & Neck Oncology, Henan Cancer Hospital, Zhengzhou, China

<sup>16</sup>Institute for Translational Epidemiology and Tisch Cancer Institute, Mount Sinai School of Medicine, New York City, NY, USA

<sup>17</sup>Department of Epidemiology, and Center for Environmental Genomics, UCLA Fielding School of Public Health, Los Angeles, CA, USA

<sup>18</sup>National Office of Cancer Prevention & Control Cancer Institute & Hospital, and Chinese Academy of Medical Sciences, Beijing, China

## Abstract

**Background**—Although the impact of oral hygiene on head and neck cancer (HNC) risk has been investigated, few studies have been conducted among Asians.

**Methods**—We conducted a multicenter case-control study to investigate this potential association. We performed unconditional multiple logistic regression models adjusted by potential confounders.

**Results**—We observed an inverse association of frequency of dental visits with HNC risk, with an adjusted OR of 3.70 (2.51-5.45) for never dental visits compared with 1 time/year ( $p_{\text{trend}} < 0.001$ ). We also observed a positive association between the number of missing teeth and HNC risk, with an adjusted OR for 5 missing teeth compared with < 5 missing teeth of 1.49 (1.08-2.04). Combining multiple oral hygiene indicators, poor oral hygiene scores increased HNC risk.

**Conclusions**—Poor oral hygiene may increase HNC risk in a Chinese population. Improving oral hygiene may contribute to reducing HNC risk in the Chinese population.

## Keywords

Oral Hygiene; Dental Care; Risk; Head and Neck cancer; Chinese

## Introduction

Head and neck cancer (HNC) is the sixth most common cancer in the world, with more than 600,000 cases diagnosed each year<sup>(1)</sup>. The anatomical sites of HNC typically include oral

cavity, oropharynx, hypopharynx and larynx, and it has been noted that these sites are strongly associated with environmental exposures. The established risk factors are predominantly tobacco smoking and alcohol drinking, and these factors cause approximately 80% of HNCs independently or synergistically<sup>(2, 3)</sup>. Recently, human papilloma virus (HPV) infection has become an established risk factor for oropharyngeal cancer<sup>(4)</sup>.

Although it has been proposed that oral hygiene and dental care indicators may modify HNC risk, it remains inconsistent. Higher proportions of missing teeth<sup>(5–13)</sup>, denture use<sup>(6, 11, 14–17)</sup> and gum bleeding<sup>(9, 13, 18)</sup> might increase HNC risk, and frequent teeth cleaning<sup>(5, 9, 10, 13, 19–21)</sup> and regular visit dentist<sup>(9–13, 16, 21, 22)</sup> might decrease HNC risk. Recently, the International Head and Neck Cancer Epidemiology Consortium (INHANCE), in a large-scale pooled case-control study, reported significant associations between oral hygiene indicators and HNC risk after adjustment of potential confounders<sup>(23)</sup>. Although this paper reported on a high impact of oral hygiene on HNC risk, only one Asian study was included in this study. Therefore, we conducted a large scale multi-center case-control study to investigate the association of oral hygiene and dental care with HNC risk in a Chinese population.

## Materials and Methods

### Design and subjects

We conducted a multicenter case-control study to elucidate the etiology of HNC in an East Asian population. The study center consists of eight centers from China (Beijing, Fujian, Henan, Jiangsu, Liaoning, Shanghai, Sichuan) and Taiwan, and the recruitment of subjects from December 2010 to February 2015. Detailed information from the face-to-face interview of both cases and controls included items on smoking and drinking habits, dietary habits, height and weight, individual and family medical history, occupational status, education level, and other lifestyle factors including oral hygiene and dental care. The study was approved by the ethical board of study centers and centrally at the University of Utah, and all subjects signed an informed consent form.

The inclusion criteria for cases were (1) age more than 18 years old, (2) confirmed invasive tumor of head and neck region (3) histologically confirmed squamous cell carcinoma, and (4) interviews performed within six months of cancer diagnosis. Tumors were assigned to one of the five categories as follows: (i) oral cavity (ICD-O-3 topography: C00.3 to C00.9, C02.0 to C02.3, C03.0, C03.1, C03.9, C04.0, C04.1, C04.8, C04.9, C05.0, C06.0 to C06.2, C06.8, and C06.9); (ii) oropharynx (C01.9, C02.4, C05.1, C05.2, C09.0, C09.1, C09.8, C09.9, C10.0, C10.2–C10.4, C10.8, and C10.9) (iii) hypopharynx (C12.9, C13.0 to C13.2, C13.8, and C13.9) (iv) oral cavity, pharynx unspecified or overlapping (C02.8, C02.9, C05.8, C05.9, C14.0, C14.2, and C14.8), and (v) larynx (C10.1, C32.0 to C32.3 and C32.8 to C32.9). Controls were frequency matched by sex, 5-year age group, ethnicity, and residence area from each of the study centers. Controls were selected from a defined list of non-chronic diseases not related to tobacco smoking or alcohol drinking. The proportion of hospital controls within a particular diagnostic group did not exceed 33%; These groups were (1) benign disorders, (2) endocrine and metabolic, (3) skin, subcutaneous tissue, and musculoskeletal disorders, (4) trauma, (5) circulatory disorders, (6) ear and eye disorders,

(7) diseases of upper-respiratory tract, (8) diseases of the oral cavity, jaw and salivary gland, (9) gastro-intestinal, (10) nervous system, (11) other diseases, and (12) no diagnosis (healthy population). Hospital controls were randomly chosen from subjects admitted as in-patients or out-patients in the same study center as the cases, and they were in the hospital for less than one month when recruited.

Finally, our study included 921 cases (424 oral cavity, 106 oropharynx, 81 hypopharynx, 85 larynx, and 225 unspecified or overlapping) and 806 controls.

### Definition of the exposure variable

Our questionnaire included four oral hygiene and dental care indicators: frequency of teeth cleaning, number of missing teeth, denture use, and frequency of dental visits. We divided subjects into each category according to oral hygiene and dental care indicators: 2 categories for frequency of teeth cleaning (> 1 time/day and ≤ 1 time/day), 2 categories for number of missing teeth (<5 and ≥ 5), 2 categories for denture use (No or Yes), and 4 categories for frequency of dental visits (≥ 1 time/year, 1 time/2-4years, 1 time/ 5 years, and never). In addition, we calculated oral hygiene scores using all oral hygiene and dental care indicators in this study. We summed the following variables: frequency of teeth cleaning: > 1 time/day=0, ≤ 1 time/day=1; number of missing teeth: '<5'=0, '≥ 5'=1; denture use: no=0, yes=1; regular dental visits: yes=0, no=1. This score ranged from 0 to 4 from best to worst oral hygiene condition.

### Statistical analysis

We estimated the odds ratios (ORs) and the 95% confidence intervals (CIs) using unconditional multiple logistic regression models. Models included adjustment for ethnicity (Han vs Taiwanese vs others), age (18-44 vs 45-54 vs 55-64 vs 65-85 years), sex (male vs female), education levels (illiterate vs primary school vs junior/middle school vs senior/high school vs college/university above), center (China mainland vs Taiwan), cigarette smoking intensity (never vs <20 cigarettes/day vs ≥ 20 cigarettes/day), cigarette smoking duration (never vs <20 years vs ≥ 20 and <40 years vs ≥ 40 years), betel quid chewing intensity (never vs <20 pieces/day vs ≥ 20 pieces/day), betel quid chewing duration (never vs <20 years vs ≥ 20 years), alcohol drinking intensity (never vs <2 drinks/day vs ≥ 2 drinks/day), alcohol drinking duration (never vs <20 years vs ≥ 20 and <40 years vs ≥ 40 years), and body mass index (BMI) at interview period (<22 kg/m<sup>2</sup> vs ≥ 22 and <25 kg/m<sup>2</sup> vs ≥ 25 kg/m<sup>2</sup>). Betel quid chewing intensity and duration were adjusted in cases with oral cavity and oropharynx, and with oral cavity, pharynx unspecified or overlapping. Differences in categorical variables across groups were assessed by the chi<sup>2</sup>-test or Fisher's exact test as appropriate. To evaluate potential interactions of oral hygiene and dental care indicators with potential confounders, we performed likelihood-ratio tests, which compared models with and without the interaction term (*p* for heterogeneity).

All statistical analyses were performed using the software STATA ver. 14 (Stata Corp, College Station, TX, USA). All tests were two-sided, and *p*-values of <0.05 were considered statistically significant.

## Results

Table 1 shows the distributions of cases and controls by subject characteristics. Among cases, the proportion of older subjects, male, cigarette smokers, betel quid consumers, alcohol drinkers, lower education level, or Han population was significantly higher than controls.

We observed a significant inverse association of frequency of dental visits with HNC risk, with adjusted ORs of 1.72 (95% CI, 1.10-2.67) for 1 time/2-4 years, 2.09 (95% CI, 1.40-3.14) for 1 time/ 5 years, and 3.70 (95% CI, 2.51-5.45) for never dental visits compared with 1 time/year with a statistically significant trend ( $p_{\text{trend}} < 0.001$ ; Table 2). In addition, the number of missing teeth was significantly associated with an increased HNC risk, with an adjusted OR for 5 missing teeth compared with < 5 missing teeth of 1.49 (95% CI, 1.08-2.04). Although lower frequency of teeth cleaning and denture use were positively associated with HNC risk, these associations were inconsistent after adjustment by potential confounders. Additionally, poor oral hygiene scores increased the risk of HNC, with adjusted ORs of 1.99 (95% CI, 1.41-2.82) for a score of 1, 1.88 (95% CI, 1.30-2.71) for a score of 2 and 4.76 (95% CI, 2.88-7.85) for a score of 3 compared with 0, with a significant trend ( $p_{\text{trend}} < 0.001$ ).

The inverse association with frequency of dental visits was significantly observed with all subsites (Table 3). The impact of frequency of teeth cleaning and number of missing teeth was stronger in oral cavity and oropharynx than in other subsites. Regarding denture use, we did not find major differences by subsite. Although poor oral hygiene scores increased the risk of all subsites, we found no significant trends for cancers risk of the hypopharynx and larynx.

We further investigated the impact of oral hygiene and dental care indicators on HNC risk stratified by potential confounders (Table 4). We observed that the impact of lower frequency of dental visits was stronger among males, smokers, drinkers and Taiwanese. Additionally, a significant inverse association with frequency of teeth cleaning was observed among Taiwanese, and we observed a significant positive association with denture use among drinkers. Regarding number of missing teeth, we observed significant associations with HNC risk among older subjects, females, never smokers, drinkers, and Taiwanese.

## Discussion

In this study, we observed that lower frequency of dental visits and greater number of missing teeth were significantly associated with an increased HNC risk in a Chinese population. In addition, we found that poor oral hygiene scores increased the risk of HNC with a significant dose-response trend.

Regarding the frequency of dental visits, to date, 12 studies have been reported<sup>(8-13, 16, 18, 21-24)</sup>. Among them, nine of them reported that lower frequency of dental visits is associated with an increased HNC risk<sup>(9-13, 16, 21-23)</sup>, and our results were consistent with them.

Next, there were 13 studies and 2 meta-analysis that evaluated the association between number of missing teeth and HNC risk<sup>(5–13, 18, 22, 23, 25–27)</sup>. Ten case-control studies and both meta-analyses supported that higher proportion of missing teeth increase HNC risk similar to our study results<sup>(5–13, 23, 26, 27)</sup>. Previous studies on missing teeth and the increased risk of HNC have been fairly consistent. The mechanism behind this association is plausible considering that periodontal disease is associated with tooth loss<sup>(28)</sup>. Zeng et al. reported on a positive association of periodontal disease with HNC risk using meta-analysis<sup>(29)</sup>. Additionally, it has been known that tooth loss is associated with smoking behavior<sup>(30)</sup>, and may be a surrogate marker of socioeconomic status (SES)<sup>(31)</sup>. Although we did not have information about the income of subjects, education level was adjusted for SES in this study.

Following the methodology from studies in Taiwan, ARCAGE, and INHANCE, we evaluated the impact of an oral hygiene score on HNC risk<sup>(13, 16, 23)</sup>. All of the previous studies, and our current study indicated a significant dose-positive relationship between oral hygiene scores and HNC risk after adjustment by smoking and drinking.

The strengths of this study includes the number of HNC cases, which is one of the largest in an Asian population. Second, we adjusted for a considerable number of potential confounders carefully including tobacco smoking, alcohol drinking and betel quid use. We were able to conduct various stratified analyses. Some limitations of our study include the hospital-based case-control design. We tried to minimize the effect of selection bias, and we selected controls with diseases unrelated to smoking and drinking. Our results could potentially be affected by overestimation of odds ratios due to recall bias, since subjects with cancer may recall poor oral hygiene with more effort. Though we would not expect that HNC patients would necessarily think of poor oral hygiene as a strong risk factor for HNC, in which case, the recall bias may be minimized. Third, residual confounding of smoking and drinking could not be ruled out completely, but we did conduct analysis among never-smokers and never-drinkers. Additionally, we adjusted on betel quid chewing intensity and duration, which is a common habit in some East Asian populations. Fourth, we did not have information on HPV infection. Although we evaluated the impact of oral hygiene and dental care indicators in oropharyngeal cases only, these results were consistent (data not shown).

In conclusion, we observed that poor oral hygiene increased the risk of HNC and its subsites in a Chinese population. Improving oral hygiene in terms of frequent dental visits may contribute to reducing HNC risk in the Chinese population.

## Acknowledgments

This investigation was supported by the University of Utah Study Design and Biostatistics Center, with funding in part from the National Cancer Institute through Cancer Center Support P30 CA042014 awarded to Huntsman Cancer Institute, and the National Center for Research Resources and the National Center for Advancing Translational Sciences, National Institutes of Health, through Grant 8UL1TR000105 (formerly UL1RR025764). In addition, this work was supported by JSPS Grant-in-Aid for Young Scientists (B) to D. Kawakita (No.15K21283). These grantors were not involved in the study design, subjects enrollment, study analysis or interpretation, or submission of the manuscript.

## References

1. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer*. 2010; 127(12):2893–917. [PubMed: 21351269]
2. Zeka A, Gore R, Kriebel D. Effects of alcohol and tobacco on aerodigestive cancer risks: a meta-regression analysis. *Cancer Causes Control*. 2003; 14(9):897–906. [PubMed: 14682447]
3. Hashibe M, Brennan P, Chuang SC, et al. Interaction between tobacco and alcohol use and the risk of head and neck cancer: pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *Cancer Epidemiol Biomarkers Prev*. 2009; 18(2):541–50. [PubMed: 19190158]
4. Gillison ML, Chaturvedi AK, Anderson WF, Fakhry C. Epidemiology of Human Papillomavirus-Positive Head and Neck Squamous Cell Carcinoma. *Journal of clinical oncology*. 2015; 33(29):3235–42. [PubMed: 26351338]
5. Zheng TZ, Boyle P, Hu HF, et al. Dentition, oral hygiene, and risk of oral cancer: a case-control study in Beijing, People's Republic of China. *Cancer Causes Control*. 1990; 1(3):235–41. [PubMed: 2102296]
6. Marshall JR, Graham S, Haughey BP, et al. Smoking, alcohol, dentition and diet in the epidemiology of oral cancer. *Eur J Cancer B Oral Oncol*. 1992; 28B(1):9–15. [PubMed: 1422474]
7. Bundgaard T, Wildt J, Frydenberg M, Elbrond O, Nielsen JE. Case-control study of squamous cell cancer of the oral cavity in Denmark. *Cancer Causes Control*. 1995; 6(1):57–67. [PubMed: 7718736]
8. Garrote LF, Herrero R, Reyes RM, et al. Risk factors for cancer of the oral cavity and oro-pharynx in Cuba. *Br J Cancer*. 2001; 85(1):46–54. [PubMed: 11437401]
9. Balaram P, Sridhar H, Rajkumar T, et al. Oral cancer in southern India: the influence of smoking, drinking, paan-chewing and oral hygiene. *Int J Cancer*. 2002; 98(3):440–5. [PubMed: 11920597]
10. Lissowska J, Pilarska A, Pilarski P, et al. Smoking, alcohol, diet, dentition and sexual practices in the epidemiology of oral cancer in Poland. *Eur J Cancer Prev*. 2003; 12(1):25–33. [PubMed: 12548107]
11. Rosenquist K, Wennerberg J, Schildt EB, Bladstrom A, Goran Hansson B, Andersson G. Oral status, oral infections and some lifestyle factors as risk factors for oral and oropharyngeal squamous cell carcinoma. A population-based case-control study in southern Sweden. *Acta Otolaryngol*. 2005; 125(12):1327–36. [PubMed: 16303683]
12. Guha N, Boffetta P, Wunsch Filho V, et al. Oral health and risk of squamous cell carcinoma of the head and neck and esophagus: results of two multicentric case-control studies. *Am J Epidemiol*. 2007; 166(10):1159–73. [PubMed: 17761691]
13. Chang JS, Lo HI, Wong TY, et al. Investigating the association between oral hygiene and head and neck cancer. *Oral Oncol*. 2013; 49(10):1010–7. [PubMed: 23948049]
14. Young TB, Ford CN, Brandenburg JH. An epidemiologic study of oral cancer in a statewide network. *Am J Otolaryngol*. 1986; 7(3):200–8. [PubMed: 3728828]
15. Piemonte ED, Lazos JP, Brunotto M. Relationship between chronic trauma of the oral mucosa, oral potentially malignant disorders and oral cancer. *J Oral Pathol Med*. 2010; 39(7):513–7. [PubMed: 20456614]
16. Ahrens W, Pohlabein H, Foraita R, et al. Oral health, dental care and mouthwash associated with upper aerodigestive tract cancer risk in Europe: the ARCAGE study. *Oral Oncol*. 2014; 50(6):616–25. [PubMed: 24680035]
17. Rotundo LD, Toporcov TN, Biazevic GH, de Carvalho MB, Kowalski LP, Antunes JL. Are recurrent denture-related sores associated with the risk of oral cancer? A case control study. *Rev Bras Epidemiol*. 2013; 16(3):705–15. [PubMed: 24896283]
18. Talamini R, Vaccarella S, Barbone F, et al. Oral hygiene, dentition, sexual habits and risk of oral cancer. *Br J Cancer*. 2000; 83(9):1238–42. [PubMed: 11027440]
19. Velly AM, Franco EL, Schlecht N, et al. Relationship between dental factors and risk of upper aerodigestive tract cancer. *Oral Oncol*. 1998; 34(4):284–91. [PubMed: 9813724]
20. Sato F, Oze I, Kawakita D, et al. Inverse association between toothbrushing and upper aerodigestive tract cancer risk in a Japanese population. *Head Neck*. 2011; 33(11):1628–37. [PubMed: 21259377]

21. Maier H, Zoller J, Herrmann A, Kreiss M, Heller WD. Dental status and oral hygiene in patients with head and neck cancer. *Otolaryngol Head Neck Surg.* 1993; 108(6):655–61. [PubMed: 8516003]
22. Divaris K, Olshan AF, Smith J, et al. Oral health and risk for head and neck squamous cell carcinoma: the Carolina Head and Neck Cancer Study. *Cancer Causes Control.* 2010; 21(4):567–75. [PubMed: 20049634]
23. Hashim D, Sartori S, Brennan P, et al. The role of oral hygiene in head and neck cancer: results from International Head and Neck Cancer Epidemiology (INHANCE) consortium. *Ann Oncol.* 2016; 27(8):1619–25. [PubMed: 27234641]
24. Eliot MN, Michaud DS, Langevin SM, McClean MD, Kelsey KT. Periodontal disease and mouthwash use are risk factors for head and neck squamous cell carcinoma. *Cancer Causes Control.* 2013; 24(7):1315–22. [PubMed: 23568534]
25. Hiraki A, Matsuo K, Suzuki T, Kawase T, Tajima K. Teeth loss and risk of cancer at 14 common sites in Japanese. *Cancer Epidemiol Biomarkers Prev.* 2008; 17(5):1222–7. [PubMed: 18483345]
26. Wang RS, Hu XY, Gu WJ, Hu Z, Wei B. Tooth loss and risk of head and neck cancer: a meta-analysis. *PLoS One.* 2013; 8(8):e71122. [PubMed: 23990929]
27. Zeng XT, Luo W, Huang W, Wang Q, Guo Y, Leng WD. Tooth loss and head and neck cancer: a meta-analysis of observational studies. *PLoS One.* 2013; 8(11):e79074. [PubMed: 24260154]
28. Chambrone L, Chambrone D, Lima LA, Chambrone LA. Predictors of tooth loss during long-term periodontal maintenance: a systematic review of observational studies. *J Clin Periodontol.* 2010; 37(7):675–84. [PubMed: 20528960]
29. Zeng XT, Deng AP, Li C, Xia LY, Niu YM, Leng WD. Periodontal disease and risk of head and neck cancer: a meta-analysis of observational studies. *PLoS One.* 2013; 8(10):e79017. [PubMed: 24194957]
30. Simila T, Virtanen JI. Association between smoking intensity and duration and tooth loss among Finnish middle-aged adults: The Northern Finland Birth Cohort 1966 Project. *BMC Public Health.* 2015; 15:1141. [PubMed: 26576994]
31. Mundt T, Polzer I, Samietz S, et al. Gender-dependent associations between socioeconomic status and tooth loss in working age people in the Study of Health in Pomerania (SHIP), Germany. *Community Dent Oral Epidemiol.* 2011; 39(5):398–408. [PubMed: 21241349]

Table 1

Characteristics of head and neck cancer cases and controls

Characteristics	Cases (N=921)		Controls (N=806)		p-value
	N	%	N	%	
<i>Age</i>					
18-44	146	16	257	32	<0.001
45-54	273	30	215	27	
55-64	297	32	222	27	
65-85	205	22	112	14	
<i>Sex</i>					
Male	726	79	556	69	<0.001
Female	195	21	250	31	
<i>Cigarette smoking intensity (cigarettes/day)</i>					
Never	319	34	462	57	<0.001
<20	148	16	136	17	
20	448	49	205	25	
Missing	6	1	3	1	
<i>Duration of cigarette smoking (years)</i>					
Never	319	35	462	57	<0.001
<20	70	8	101	13	
20 and <40	391	42	192	24	
40	138	15	49	6	
Missing	3	0	2	0	
<i>Betel quid chewing intensity (betel pieces/day)</i>					
Never	624	68	761	94	<0.001
<20	113	12	28	4	
20	166	18	14	2	
Missing	18	2	3	0	
<i>Duration of betel quid chewing (years)</i>					
Never	624	68	761	95	<0.001
<20	117	13	31	4	

Characteristics	Cases (N=921)		Controls (N=806)		p-value
	N	%	N	%	
20	175	19	11	1	
Missing	5	0	3	0	
<b>Alcohol drinking intensity (drinks/day)</b>					
Never	433	47	582	72	<0.001
<2	149	16	133	17	
2	297	32	72	9	
Missing	42	5	19	2	
<b>Duration of drinking (years)</b>					
Never	433	47	582	72	<0.001
<20	106	11	93	11	
20 and <40	272	30	104	13	
40	77	8	22	3	
Missing	33	4	5	1	
<b>BMI (kg/m<sup>2</sup>)</b>					
<22	306	33	241	30	0.332
22 and <25	329	36	300	37	
>25	285	31	264	33	
Missing	1	0	1	0	
<b>Education</b>					
Illiterate	59	6	24	3	<0.001
Primary school	228	25	129	16	
Junior/middle school	261	28	150	19	
Senior/high school	244	27	170	21	
College/university and above	129	14	333	41	
<b>Center</b>					
Mainland	439	48	405	50	0.284
Taiwan	482	52	401	50	
<b>Ethnicity</b>					
Han	556	60	407	51	<0.001
Taiwanese	348	38	390	48	

Characteristics	Cases (N=921)		Controls (N=806)		p-value
	N	%	N	%	
Others	17	2	9	1	
<i>Subsite</i>					
Oral cavity	424	46			
Oropharynx	106	12			
Hypopharynx	81	9			
Larynx	85	9			
Unspecified or overlapping	225	24			

Abbreviation; BMI: body mass index.

**Table 2**

Impact of oral hygiene and dental care on head and neck cancer risk

Variables	Case	Control	OR	95% CI	OR**	95% CI
<i>Frequency of teeth cleaning</i>						
>1 time/day	429	491	1.00	-	1.00	-
1 time/day	484	313	1.77	1.46-2.15	1.13	0.89-1.44
Missing	8	2	-	-	-	-
<i>Number of missing teeth</i>						
<5	635	696	1.00	-	1.00	-
5	264	108	2.68	2.09-3.43	1.49	1.08-2.04
Missing	22	2	-	-	-	-
<i>Denture use</i>						
No	719	689	1.00	-	1.00	-
Yes	197	114	1.66	1.29-2.13	1.29	0.94-1.75
Missing	5	3	-	-	-	-
<i>Frequency of dental visits</i>						
1 time/year	89	271	1.00	-	1.00	-
1 time/2-4 years	88	115	2.33	1.61-3.36	1.72	1.10-2.67
1 time/ 5 years	205	132	4.73	3.42-6.54	2.09	1.40-3.14
Never	526	261	6.14	4.63-8.13	3.70	2.51-5.45
Missing	13	27	-	-	-	-
<i>P<sub>trend</sub></i>				<0.001		<0.001
<i>Oral hygiene score (best to worst oral hygiene)</i>						
0	124	288	1.00	-	1.00	-
1	290	249	2.71	2.06-3.54	1.99	1.41-2.82
2	305	202	3.51	2.66-4.62	1.88	1.30-2.71
3	163	37	10.23	6.76-15.49	4.76	2.88-7.85
Missing	39	30	-	-	-	-
<i>P<sub>trend</sub></i>				<0.001		<0.001

Abbreviation; OR, odds ratio; CI, confidence interval.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

\* Oral hygiene score is the sum of the four following variables: frequency of teeth cleaning: >1 time/day=0, 1 time/day=1; number of missing teeth: <5=0, 5=1; denture use: no=0, yes=1; regular dental visits: yes=0, no=1.

\*\* Adjusted by age, sex, cigarette smoking intensity, cigarette smoking duration, betel quid chewing intensity, betel quid chewing duration, drinking intensity, drinking duration, body mass index, study center, ethnicity, education level.

**Table 3**

Impact of oral hygiene and dental care on head and neck cancer risk according to subsites and study center

Variables	<u>Oral cavity and Oropharynx</u>		<u>Hypopharynx and Larynx</u>		<u>Unspecified or Overlapping</u>	
	OR**	95% CI	OR**	95% CI	OR**	95% CI
<b>Frequency of teeth cleaning</b>						
>1 time/day	1.00	-	1.00	-	1.00	-
1 time/day	1.49	1.11-1.99	0.66	0.42-1.05	1.14	0.80-1.64
<b>Number of missing teeth</b>						
<5	1.00	-	1.00	-	1.00	-
5	1.83	1.27-2.64	1.30	0.74-2.27	1.35	0.84-2.18
<b>Denture use</b>						
No	1.00	-	1.00	-	1.00	-
Yes	1.34	0.94-1.91	1.12	0.62-2.02	1.35	0.83-2.17
<b>Frequency of dental visits</b>						
1 time/year	1.00	-	1.00	-	1.00	-
1 time/2-4 years	2.13	1.21-3.74	1.06	0.34-3.28	1.68	0.89-3.18
1 time/ 5 years	1.75	1.03-2.97	2.82	1.19-6.70	2.45	1.38-4.33
Never	3.68	2.24-6.05	6.42	2.77-14.88	4.34	2.43-7.76
<i>P<sub>trend</sub></i>	<0.001		<0.001		<0.001	
<b>Oral hygiene score (best to worst oral hygiene)</b>						
0	1.00	-	1.00	-	1.00	-
1	1.79	1.14-2.82	4.96	2.19-11.25	2.48	1.52-4.02
2	2.29	1.44-3.63	2.65	1.14-6.14	1.94	1.13-3.33
3	6.30	3.51-11.29	6.23	2.30-16.90	5.06	2.43-10.54
<i>P<sub>trend</sub></i>	<0.001		0.106		<0.001	

Abbreviation; OR, odds ratio; CI, confidence interval.

\* Oral hygiene score is the sum of the four following variables: frequency of teeth cleaning: >1 time/day=0, 1 time/day=1; number of missing teeth: <5=0, 5=1; denture use: no=0, yes=1; regular dental visits: yes=0, no=1.

\*\* Adjusted by age, sex, cigarette smoking intensity, cigarette smoking duration, betel quid chewing intensity, betel quid chewing duration, drinking intensity, drinking duration, body mass index, study center, ethnicity, education level. Betel quid chewing intensity and betel quid chewing duration was not adjusted in hypopharynx and larynx.

**Table 4**

Impact of oral hygiene and dental care on head and neck cancer risk according to potential confounders

Variables	Age				Sex				Smoking status				Drinking status				Study center			
	18-54 years		55-85 years		Male		Female		Never		Ever		Never		Ever		Main land		Taiwan	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
<b>Frequency of teeth cleaning</b>																				
>1 time/day	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-
1 time/day	1.48	1.05-2.09	0.92	0.65-1.31	1.17	0.88-1.57	1.20	0.76-1.91	1.05	0.74-1.49	1.24	0.88-1.75	1.34	0.99-1.82	0.87	0.58-1.31	0.94	0.69-1.29	1.79	1.18-2.71
$P_{\text{for heterogeneity}}$	0.129		0.650		0.420		0.046												0.029	
<b>Number of missing teeth</b>																				
<5	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-
5	1.39	0.72-2.72	1.66	1.16-2.37	1.36	0.93-1.99	1.91	1.04-3.52	2.05	1.28-3.28	1.09	0.70-1.68	1.30	0.88-1.93	2.09	1.18-3.72	1.41	0.90-2.21	1.68	1.03-2.71
$P_{\text{for heterogeneity}}$	0.792		0.107		0.009		0.207												0.726	
<b>Denture use</b>																				
No	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-
Yes	1.49	0.80-2.79	1.30	0.91-1.86	1.18	0.81-1.73	1.42	0.82-2.45	1.37	0.89-2.11	1.18	0.75-1.84	0.97	0.66-1.44	2.14	1.23-3.71	1.36	0.93-1.99	1.19	0.67-2.14
$P_{\text{for heterogeneity}}$	0.708		0.338		0.372		0.017												0.244	
<b>Frequency of dental visits</b>																				
1 time/year	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-
1 time/2-4 years	2.23	1.18-4.23	1.57	0.85-2.91	2.70	1.55-4.70	0.78	0.36-1.71	1.15	0.62-2.11	2.80	1.45-5.39	1.63	0.98-2.71	1.93	0.80-4.67	0.65	0.30-1.41	2.57	1.43-4.62
1 time/ 5 years	2.91	1.59-5.35	1.87	1.07-3.26	3.69	2.23-6.11	0.76	0.36-1.60	1.48	0.83-2.62	3.04	1.68-5.51	1.62	0.98-2.69	3.10	1.47-6.52	0.62	0.30-1.29	4.10	2.43-6.94
Never	4.41	2.43-8.02	3.80	2.23-6.46	6.85	4.16-11.26	1.27	0.65-2.46	2.11	1.24-3.60	6.70	3.73-12.03	3.02	1.90-4.81	5.58	2.67-11.65	1.05	0.55-2.01	13.30	7.11-24.89
$P_{\text{for heterogeneity}}$	0.434		<0.001		0.003		0.130												<0.001	
<b>Oral hygiene score (best to worst oral hygiene)</b>																				
0	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-	1.00	-
1	2.48	1.56-3.94	1.66	0.96-2.85	2.93	1.89-4.54	1.00	0.54-1.84	1.27	0.79-2.05	3.36	1.99-5.65	1.70	1.13-2.57	3.32	1.69-6.50	1.43	0.80-2.56	2.95	1.84-4.75
2	2.67	1.57-4.53	1.56	0.91-2.66	2.86	1.80-4.54	0.90	0.47-1.73	1.28	0.76-2.13	2.89	1.68-4.97	1.74	1.12-2.73	2.72	1.38-5.37	1.33	0.75-2.37	3.05	1.71-5.42
3	5.68	2.22-14.51	4.30	2.28-8.13	5.06	2.75-9.30	6.14	2.28-16.49	3.92	1.96-7.83	5.97	2.86-12.48	3.69	2.01-6.77	####	4.02-27.77	3.01	1.47-6.15	11.06	4.47-27.39
$P_{\text{for heterogeneity}}$	0.579		0.185		0.367		0.323												0.159	

Abbreviation; OR, odds ratio; CI, confidence interval.

\* Oral hygiene score is the sum of the four following variables: frequency of teeth cleaning: 1 time/day=0, <1 time/day=1; number of missing teeth: <5=0, 5=1; denture use: no=0, yes=1; regular dental visits: yes=0, no=1. Adjusted by age, sex, cigarette smoking intensity, cigarette smoking duration, betel quid chewing intensity, betel quid chewing duration, drinking intensity, drinking duration, body mass index, study center, ethnicity, education level.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript